

CGPG-94/4-6
April 1994

Real formulations of complex gravity and a complex formulation of real gravity

Peter Peldán ¹

Center for Gravitational Physics and Geometry
The Pennsylvania State University, University Park, PA 16802, USA

Abstract

Two gauge and diffeomorphism invariant theories on the Yang-Mills phase space are studied. They are based on the Lie-algebras $so(1, 3)$ and $\widetilde{so(3)}$ – the loop-algebra of $so(3)$. Although the theories are manifestly real, they can both be reformulated to show that they describe complex gravity and an infinite number of copies of complex gravity, respectively. The connection to real gravity is given. For these theories, the reality conditions in the conventional Ashtekar formulation are represented by normal constraint-like terms.

PACS number: 04.20.Fy

¹Email address: peldan@phys.psu.edu

1 Introduction

In the search for a generalization of the Ashtekar variables [1] to other gauge groups, it has previously been shown [2], [3], [4] that there exist an arbitrary gauge group generalization in both (2+1) as well as (3+1)-dimensions. Moreover, for both these cases, the generalized theories were shown to produce the conventional Yang-Mills theory when expanded around de Sitter spacetime. However, the problem with the (3+1)-dimensional model is that the theory is required to be complex in order to include Lorentzian spacetimes, and no reality conditions have yet been found. Another expectation one may have on a unified description of gravity and Yang-Mills theory, is that the theory itself should be somewhat restrictive regarding what gauge groups that could be used. This is not the case for the models presented in [2] and [4], these constructions allow all gauge groups whose Lie-algebras admit a non-degenerate invariant bilinear form. Thus, motivated by the desire to solve these two problems, I have, in this paper, studied another type of generalization of the Ashtekar formulation. The method used in [4] was to replace the structure constant of $so(3)$ by the three-dimensional internal Levi-Civita symbol, which then was generalized to be a well defined object for higher dimensional Lie-algebras. In this paper, I instead keep the structure constants in the theory, and use only Lie-algebras whose structure constants satisfy an identity of the type

$$f_{IJK}f^K{}_{LM} \sim \sum_{i,j} g^i_{I[L}g^j_{M]J} \quad (1.1)$$

where the g^i_{IJ} are invariant bilinear forms of the Lie-algebra. I show that it is possible to find a gauge and diffeomorphism invariant theory based on two such Lie-algebras: $so(1,3)$ and the loop-algebra $\widetilde{so(3)}$. However, as can be shown by reformulating the theories, both these models describe nothing more than complex gravity. The $so(1,3)$ theory is exactly equivalent to complex gravity, while the $\widetilde{so(3)}$ theory corresponds to an infinite number of copies of complex gravity. The interesting fact about these formulations is perhaps that they represent a way of getting real formulations of complex gravity, and they may possibly be used to shed some light over the reality condition-problem in Ashtekar's variables.

The models presented here are constructed in (3+1)-dimensions for the Lie-algebras $so(1,3)$ and $\widetilde{so(3)}$. But it is straightforward to generalize the formulation to other dimensions $\geq (2+1)$ and for the Lie-algebras $so(4)$, $so(2,2)$, $so(1,2)$ – and isomorphic ones – as well as their loop-algebras.

2 What is the problem with arbitrary gauge groups?

The problem of using arbitrary gauge groups for the Ashtekar Hamiltonian comes from the constraint algebra; the algebra fails to close in general. In the Ashtekar formulation, the first class constraints are

$$\mathcal{G}^I := \mathcal{D}_a E^{aI} = \partial_a E^{aI} + f^I{}_{JK} A_a^J E^{aK} \approx 0 \quad (2.1)$$

$$\mathcal{H}_a := \frac{1}{2} \epsilon_{abc} E^{bI} B^c_I \approx 0 \quad (2.2)$$

$$\mathcal{H} := \frac{1}{4} \epsilon_{abc} f_{IJK} E^{aI} E^{bJ} B^{cK} \approx 0 \quad (2.3)$$

They are called, the Gauss law, the vector constraint, and the Hamiltonian constraint, respectively. The conventions are: $a, b, c, \dots$ are spatial indices on the hypersurface, $I, J, K, \dots$ are Lie-algebra indices in the vector representation. The structure constant is denoted f_{IJK} , and Lie-algebra indices are raised and lowered by an invariant bilinear form on the Lie-algebra. (For the conventional Ashtekar Hamiltonian, which together with a set of reality conditions describes real Einstein gravity, the Lie-algebra is $so(3; C)$ and the bilinear form is δ_{IJ} .) The basic conjugate fields are a gauge connection A_{aI} and "the electric field" E^{aI} . They satisfy: $\{A_{aI}(x), E^{bJ}(y)\} = \delta_I^J \delta_a^b \delta^3(x - y)$. B^{aI} is the "magnetic field": $B^{aI} := \epsilon^{abc} F_{bc}^I = \epsilon^{abc} (2\partial_b A_c^I + f_{JK}^I A_b^J A_c^K)$. The totally antisymmetric Levi-Civita tensor densities are ϵ^{abc} and ϵ_{abc} . They are taken to have $\epsilon^{123} = \epsilon_{123} = 1$ in every coordinate system.

Since $\mathcal{G}^I$ and $\mathcal{H}_a := \mathcal{H}_a - A_{aI} \mathcal{G}^I$ easily can be shown to generate gauge transformations and spatial diffeomorphisms, there can be no problems with the Poisson brackets containing these constraints, as long as all constraints are gauge- and diffeomorphism covariant. The only potential obstruction to get a closed constraint algebra, thus comes from $\{\mathcal{H}[N], \mathcal{H}[M]\}$. A straightforward calculation gives

$$\{\mathcal{H}[N], \mathcal{H}[M]\} = \frac{1}{2} \int_{\Sigma} d^3x \left((N\partial_a M - M\partial_a N) E^{aI} E^{bJ} \epsilon_{bcd} E^{cK} B^{dL} f_{IJM} f^M{}_{KL} \right) \quad (2.4)$$

A right-hand side which in general is *not* a linear combination of constraints. For $SO(3)$ or locally isomorphic groups, the structure constant equals the epsilon symbol, and therefore the right-hand side above reduces to the vector constraint smeared over some test function. For arbitrary gauge groups, no such simplification occurs, and the constraint algebra therefore fails to close. Thus, it seems that the requirement on the Lie-algebra is that its structure constants must satisfy $f_{IJ}^K f_{KLM} \sim g_{IL} g_{JM} - g_{IM} g_{JL}$ in order to make the constraints in eq. (2.1)-(2.3) form a first class set. Here, g_{IJ} is an invariant bilinear form on the Lie-algebra. This is a severe restriction on the Lie-algebra, and in fact I only know of two algebras that satisfy this: $so(3)$ and $so(1, 2)$. However, in this paper, I will show how one can use a slightly weaker condition: suppose we have an Lie-algebra that satisfies

$$f_{IJK} f^K{}_{LM} \sim \sum_{i,j} g_{[L}^i g_{M]J}^j \quad (2.5)$$

where g_{IJ}^i are invariant bilinear forms of the Lie-algebra. The indices i and j just label the different bilinear forms. Using this in (2.4) above, we see that the right hand side becomes a linear combination of the vector constraint and "vector constraint-like" terms. By introducing the "vector constraint-like" terms as new constraints in the theory, and continue to check the algebra, one soon notices that the new constraints also produce new "Hamiltonian constraint-like" terms, which also have to be included in the set of constraints. However, if the Lie-algebra is such that the product of the bilinear forms also is invariant, the procedure stops here, and we have a first class set. That is, I require

$$g_{IJ}^i g^{jJ}{}_K = \sum_k a_k g_{IK}^k \quad (2.6)$$

where I have selected one of the bilinear forms, say g_{IJ}^0 , and its inverse to raise and lower Lie-algebra indices.

I do not know of any general theorems concerning what kind of Lie-algebras that satisfy these requirements. In the following sections, two different examples of this construction will be presented. They are based on $so(1, 3)$ and the loop-algebra $\widetilde{so(3)}$.

3 The $SO(1,3)$ theory

In this section, I will present an Ashtekar-like theory based on the Lie-algebra $so(1,3)$. The construction also works for the algebras $so(4)$ and $so(2,2)$. First I describe the Lie-algebra in subsection 3.1. In subsection 3.2, the Hamiltonian is presented and a constraint analysis is performed. In subsection 3.3, the theory is split up into self dual and antiself dual parts, and I show that the real $so(1,3)$ theory is equivalent to the complex $so(3)$ Ashtekar formulation. Finally, in subsection 3.4, two manifestly covariant and real Lagrangians for the $so(1,3)$ theory are derived.

3.1 The $so(1,3)$ algebra

Before introducing the theory, I will give a short description of the Lie-algebra. First, a basis is chosen for the fundamental representation:

$$\begin{aligned} T_{1A}{}^B &= \begin{pmatrix} 0 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix} & T_{2A}{}^B &= \begin{pmatrix} 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix} & T_{3A}{}^B &= \begin{pmatrix} 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 \end{pmatrix} \\ T_{4A}{}^B &= \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & -1 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix} & T_{5A}{}^B &= \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & -1 \\ 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \end{pmatrix} & T_{6A}{}^B &= \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & -1 & 0 \end{pmatrix} \end{aligned} \quad (3.1)$$

where $A, B, \dots$ are four-dimensional Lorentz-indices. I will also use $I, J, K, \dots$ as six-dimensional indices in the vector representation. T_1, T_2 and T_3 are the boost-generators, and T_4, T_5, T_6 are the generators of rotations. Using this representation, it is straightforward to calculate the commutator algebra of these basis elements:

$$[T_I, T_J] = f_{IJ}{}^K T_K \quad (3.2)$$

where

$$\begin{aligned} f_{12}{}^4 &= f_{31}{}^5 = f_{23}{}^6 = f_{45}{}^6 = f_{64}{}^5 = f_{56}{}^4 = f_{14}{}^2 = f_{51}{}^3 = f_{42}{}^1 = f_{26}{}^3 = f_{35}{}^1 = f_{63}{}^2 = 1 \\ f_{16}{}^i &= f_{25}{}^i = f_{34}{}^i = 0 \end{aligned} \quad (3.3)$$

Now, I need to introduce a non-degenerate invariant bilinear form – or a scalar product – in the Lie-algebra. Invariance here means, invariance under conjugation (gauge transformations). If the bilinear form is denoted $\langle A, B \rangle$ for A and B belonging to the Lie-algebra, the invariance requirement becomes:

$$\langle [C, A], B \rangle + \langle A, [C, B] \rangle = 0 \quad (3.4)$$

for all elements A, B and C belonging to the Lie-algebra. Or, expressing A, B and C in the basis T_I above: $A = A^I T_I$ etc., the invariance condition reads

$$f_{IJ}{}^K g_{KL} + f_{IL}{}^K g_{KJ} = 0 \quad (3.5)$$

where I have defined $g_{IJ} := \langle T_I, T_J \rangle$. With the f_{IJ}^K 's given above, it is easy to solve this equation for g_{IJ} , in this representation. The solution is that there exist only two different non-degenerate invariant bilinear forms

$$g_{IJ} = \begin{pmatrix} -1 & 0 & 0 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 & 0 & 0 \\ 0 & 0 & -1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix} \quad g_{IJ}^* = \begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 \end{pmatrix} \quad (3.6)$$

Of course, any linear combination of these matrices will also be an invariant bilinear form. The reason why this Lie-algebra allows two different "group-metrics" can be traced back to the fact that it splits into two subalgebras $so(1, 3; C) = so(3, C) \oplus so(3, C)$. If one wants an explicit definition of g_{IJ} it is simply given by the Cartan-Killing form $-\frac{1}{4}f_{IK}^L f_{JL}^K$ or the matrix-trace $-\frac{1}{2}Tr(T_I T_J)$. The explicit definition of g_{IJ}^* can be found by first defining the dual basis $T_{IA}^{*B} := \frac{1}{2}\epsilon_A^{BCD} T_{ICD}$ and then take the matrix-trace $-\frac{1}{2}Tr(T_I T_J^*)$. Here, I have introduced the totally antisymmetric ϵ^{ABCD} and the Lorentz-indices are raised and lowered with the Minkowski metric $\eta_{AB} = diag(-1, 1, 1, 1)$. Thus, in the fundamental representation, g_{IJ} corresponds to "tracing" with two Minkowski metrics, while g_{IJ}^* represents the ϵ_{ABCD} -trace.

Now, what is interesting about this Lie-algebra, for our purposes, is that its structure constants satisfy the identity

$$f_{IJ}^K f_{LK}^M = \delta_{[I}^M g_{J]L} + g_{[J}^*{}^M g_{I]L}^* \quad (3.7)$$

where I have defined $g_I^{*J} := g^{JK} g_{IK}^*$ and g^{IJ} is the inverse of g_{IJ} . From now on I will raise and lower Lie-algebra indices with g_{IJ} and g^{IJ} . If an index is lowered with g_{IJ}^* , the new object will be denoted with an extra *: *e.g.* $f_{IJK}^* := f_{IJ}^L g_{KL}^*$. Note that the identity (3.7) is exactly of the type (2.5), and that the product of two bilinear forms always will be an invariant bilinear form. Now, I summarize all the important definitions and relations for this algebra:

$$\begin{aligned} g_{IJ} &:= -\frac{1}{2}Tr(T_I T_J) = -\frac{1}{2}T_{IA}^B T_{JB}^A & g_{IJ}^* &:= -\frac{1}{2}Tr(T_I T_J^*) = -\frac{1}{4}\epsilon^{ABCD} T_{ICD} T_{JBA} \\ g^{IJ} g_{JK} &= \delta_K^I & g^{*IJ} &:= g^{IK} g^{*JL} g_{KL}^* \\ g^{*IJ} g_{JK}^* &= -\delta_K^I & [T_I, T_J] &= f_{IJ}^K T_K \\ f_{IJL} &:= f_{IJ}^K g_{KL} = -Tr(T_I T_J T_L) & f_{IJK}^* &:= f_{IJ}^L g_{KL}^* \\ f_{IJK}^{**} &= -f_{IJK} & f_{IJ}^K f_{LK}^M &= \delta_{[I}^M g_{J]L} + g_{[J}^*{}^M g_{I]L}^* \end{aligned} \quad (3.8)$$

3.2 The $so(1, 3)$ Hamiltonian

Then, it is time to start building a gauge and diffeomorphism invariant theory based on this Lie-algebra. The fundamental phase space variables are chosen to be a connection A_{aI} and its conjugate momenta E^{aI} :

$$\{A_{aI}(x), E^{bJ}(y)\} = \delta_a^b \delta_I^J \delta^3(x - y) \quad (3.9)$$

. More generally, one could have chosen the right-hand side to be $\delta_a^b(\delta_I^J + ag_I^{*J})$. However, this Poisson bracket may be brought to the standard one (3.9) by a simple field-redefinition: $\tilde{E}^{aI} := \frac{1}{1+a^2}(\delta_I^J - ag_J^{*I})E^{aJ}$. Now, according to the outline of the constraint algebra calculation, given in section 2, we know that the full set of constraints we need includes the conventional Ashtekar constraints as well as the new constraints that are produced by replacing δ_J^I by g_J^{*I} in the vector and the Hamiltonian constraint. Thus, we have

$$\mathcal{G}_I := \mathcal{D}_a E_I^a = \partial_a E_I^a + f_{IJK} A_a^J E^{aK} \approx 0 \quad (3.10)$$

$$\mathcal{H}_a := \frac{1}{2} \epsilon_{abc} E^{bI} B_I^c \approx 0 \quad (3.11)$$

$$\mathcal{H}_a^* := \frac{1}{2} \epsilon_{abc} E^{bI} B_I^{*c} \approx 0 \quad (3.12)$$

$$\mathcal{H} := \frac{1}{4} \epsilon_{abc} f_{IJK} E^{aI} E^{bJ} B^{cK} + \frac{\Lambda}{6} \epsilon_{abc} f_{IJK}^* E^{aI} E^{bJ} E^{cK} \approx 0 \quad (3.13)$$

$$\mathcal{H}^* := \frac{1}{4} \epsilon_{abc} f_{IJK}^* E^{aI} E^{bJ} B^{cK} - \frac{\Lambda}{6} \epsilon_{abc} f_{IJK} E^{aI} E^{bJ} E^{cK} \approx 0 \quad (3.14)$$

Here, I have introduced a cosmological constant term as well, and the conventions are chosen such that this theory will contain the conventional Ashtekar formulation for a real Λ .

The claim is now that these constraints form a first class set, and to prove that, one simply calculates the constraint algebra. First, to simplify the calculations, one may note that $\mathcal{G}^I$, $\tilde{\mathcal{H}}_a := \mathcal{H}_a - A_{aI} \mathcal{G}^I$ and $\tilde{\mathcal{H}}_a^* := \mathcal{H}_a^* - A_{aI}^* \mathcal{G}^I$ generate gauge transformations, spatial diffeomorphisms and spatial diffeomorphisms times a duality rotation, respectively:

$$\delta^{\mathcal{G}^I} E^{aI} := \{E^{aI}, \mathcal{G}^J[\Lambda_J]\} = f^I{}_{JK} \Lambda^J E^{aK} \quad (3.15)$$

$$\delta^{\mathcal{G}^I} A_a^I := \{A_a^I, \mathcal{G}^J[\Lambda_J]\} = -\mathcal{D}_a \Lambda^I \quad (3.16)$$

$$\delta^{\tilde{\mathcal{H}}_a} E^{aI} := \{E^{aI}, \tilde{\mathcal{H}}_b[N^b]\} = N^b \partial_b E^{aI} - E^{bI} \partial_b N^a + E^{aI} \partial_b N^b = \mathcal{L}_{N^b} E^{aI} \quad (3.17)$$

$$\delta^{\tilde{\mathcal{H}}_a} A_a^I := \{A_a^I, \tilde{\mathcal{H}}_b[N^b]\} = N^b \partial_b A_a^I + A_b^I \partial_a N^b = \mathcal{L}_{N^b} A_a^I \quad (3.18)$$

$$\delta^{\tilde{\mathcal{H}}_a^*} E^{aI} := \{E^{aI}, \tilde{\mathcal{H}}_b^*[N^b]\} = N^b \partial_b E^{*aI} - E^{*bI} \partial_b N^a + E^{*aI} \partial_b N^b = \mathcal{L}_{N^b} E^{*aI} \quad (3.19)$$

$$\delta^{\tilde{\mathcal{H}}_a^*} A_a^I := \{A_a^I, \tilde{\mathcal{H}}_b^*[N^b]\} = N^b \partial_b A_a^{*I} + A_b^{*I} \partial_a N^b = \mathcal{L}_{N^b} A_a^{*I} \quad (3.20)$$

With this knowledge at hand, it is a simple task to calculate all Poisson brackets containing these constraints. Since $\mathcal{G}^I$ is gauge covariant, $\mathcal{H}$ and $\mathcal{H}^*$ are gauge invariant, and all constraints are diffeomorphism covariant, we get

$$\{\mathcal{G}^I[\Lambda_I], \mathcal{G}^J[\Gamma_J]\} = \mathcal{G}^K[f_{KIJ} \Lambda^I \Gamma^J] \quad (3.21)$$

$$\{\mathcal{G}^I[\Lambda_I], \tilde{\mathcal{H}}_a[N^a]\} = \mathcal{G}^I[-\mathcal{L}_{N^a} \Lambda^I] \quad (3.22)$$

$$\{\mathcal{G}^I[\Lambda_I], \tilde{\mathcal{H}}_a^*[N^a]\} = \mathcal{G}^I[-\mathcal{L}_{N^a} \Lambda^{*I}] \quad (3.23)$$

$$\{\mathcal{G}^I[\Lambda_I], \mathcal{H}[N]\} = 0 \quad (3.24)$$

$$\{\mathcal{G}^I[\Lambda_I], \mathcal{H}^*[N]\} = 0 \quad (3.25)$$

$$\{\tilde{\mathcal{H}}_a[N^a], \tilde{\mathcal{H}}_b[M^b]\} = \tilde{\mathcal{H}}_a[-\mathcal{L}_{M^b} N^a] \quad (3.26)$$

$$\{\tilde{\mathcal{H}}_a[N^a], \tilde{\mathcal{H}}_b^*[M^b]\} = \tilde{\mathcal{H}}_a^*[-\mathcal{L}_{M^b} N^a] \quad (3.27)$$

$$\{\tilde{\mathcal{H}}_a^*[N^a], \tilde{\mathcal{H}}_b[M^b]\} = \tilde{\mathcal{H}}_a[\mathcal{L}_{M^b} N^a] \quad (3.28)$$

$$\{\mathcal{H}[N], \tilde{\mathcal{H}}_a[N^a]\} = \mathcal{H}[-\mathcal{L}_{N^a} N] \quad (3.29)$$

$$\{\mathcal{H}[N], \tilde{\mathcal{H}}_a^*[N^a]\} = \mathcal{H}^*[-\mathcal{L}_{N^a}N] \quad (3.30)$$

$$\{\mathcal{H}^*[N], \tilde{\mathcal{H}}_a[N^a]\} = \mathcal{H}^*[-\mathcal{L}_{N^a}N] \quad (3.31)$$

$$\{\mathcal{H}^*[N], \tilde{\mathcal{H}}_a^*[N^a]\} = \mathcal{H}[\mathcal{L}_{N^a}N] \quad (3.32)$$

The remaining Poisson brackets are those that only contains $\mathcal{H}$ and $\mathcal{H}^*$. A straightforward calculation gives

$$\begin{aligned} \{\mathcal{H}[N], \mathcal{H}[M]\} &= \mathcal{H}_b[E^{bI}E_I^a(M\partial_aN - N\partial_aM)] - \\ &\quad \mathcal{H}_b^*[E^{*bI}E_I^a(M\partial_aN - N\partial_aM)] \end{aligned} \quad (3.33)$$

$$\begin{aligned} \{\mathcal{H}[N], \mathcal{H}^*[M]\} &= \mathcal{H}_b[E^{bI}E_I^{*a}(M\partial_aN - N\partial_aM)] + \\ &\quad \mathcal{H}_b^*[E^{bI}E_I^a(M\partial_aN - N\partial_aM)] \end{aligned} \quad (3.34)$$

$$\begin{aligned} \{\mathcal{H}^*[N], \mathcal{H}^*[M]\} &= -\mathcal{H}_b[E^{bI}E_I^a(M\partial_aN - N\partial_aM)] + \\ &\quad \mathcal{H}_b^*[E^{*bI}E_I^a(M\partial_aN - N\partial_aM)] \end{aligned} \quad (3.35)$$

where I had to use the structure constant identity (3.7). Thus, the constraint algebra closes, and the total Hamiltonian

$$H^{tot} := \int_{\Sigma} d^3x \left(N\mathcal{H} + N^*\mathcal{H}^* + N^a\mathcal{H}_a + N^{*a}\mathcal{H}_a^* - A_{0I}\mathcal{G}^I \right) \quad (3.36)$$

defines a consistent theory. Consistent, in the sense that a field configuration that starts out on the constraint surface stays there under time evolution. (The $*$'s on the Lagrange multiplier fields are just for notational convenience.)

Given this theory, there are of course several question to ask: *e.g.* 1. What is the interpretation of the theory, what does it describe? 2. Is this theory diffeomorphism invariant (in the four-dimensional sense). 3. Is there a geometrical interpretation of the fields? And, in that case, what kind of geometry can the theory describe? Lorentzian, Euclidean, ...? 4. What is the physical meaning of the local symmetry generated by $\mathcal{H}_a^*$ and $\mathcal{H}^*$; What are the finite symmetry transformations? The answers to all these questions will be found from the result of the split of the theory in the next subsection.

3.3 $so(1, 3; C) = so(3; C) \oplus so(3; C)$

In this subsection, I will use the fact that the Lorentz Lie-algebra splits into two copies of complex $so(3)$ algebras, to decompose the $so(1, 3)$ theory. This splitting is performed by introducing self dual and antiself dual fields, and since the square of the duality-operation, in Minkowskian spacetime, equals -1, self duality necessarily means introducing complex numbers. If we instead would use the Lie-algebras $so(4)$ or $so(2, 2)$, we could still get by with real fields, and the final result would be completely different.

In the vector notation, introduced in the previous subsection, self duality and antiself duality mean;

$$q_I^{*J}V_J^{(\pm)} = \pm iV_I^{(\pm)} \quad (3.37)$$

for a self dual field $V_I^{(+)}$ and antiself dual field $V_I^{(-)}$. Given any field V_I that take values in the Lie-algebra $so(1, 3)$ – in the vector representation – the (anti)self dual part is defined as follows

$$V_I^{(\pm)} := \delta_I^{(\pm)J}V_J = \frac{1}{2}(\delta_I^J \mp ig_I^{*J})V_J \quad (3.38)$$

From this it is obvious that $V_I = V_I^{(+)} + V_I^{(-)}$, and that the dual of V_I is $V_I^* := g_I^{*J} V_J = i(V_I^{(+)} - V_I^{(-)})$. The (anti)self dual projection operator has the following features:

$$\delta_I^{(+J)} \delta_J^{(+K)} = \delta_I^{(+K)} \quad \delta_I^{(+J)} \delta_J^{(-K)} = 0 \quad \delta_I^{(-J)} \delta_J^{(-K)} = \delta_I^{(-K)} \quad (3.39)$$

Furthermore, since these projection operators are built out of invariant forms of the $so(1,3)$ algebra, it follows that

$$\delta_I^{(+J)} f_{JKL} = \delta_I^{(+J)} \delta_K^{(+M)} f_{JML} = \delta_I^{(+J)} \delta_K^{(+M)} \delta_L^{(+N)} f_{JMN} := f_{IKL}^{(+)} \quad (3.40)$$

$$\delta_I^{(-J)} f_{JKL} = \delta_I^{(-J)} \delta_K^{(-M)} f_{JML} = \delta_I^{(-J)} \delta_K^{(-M)} \delta_L^{(-N)} f_{JMN} := f_{IKL}^{(-)} \quad (3.41)$$

$$\delta_I^{(+J)} \delta_K^{(-M)} f_{JML} = 0 \quad (3.42)$$

$$f_{IJK} = f_{IJK}^{(+)} + f_{IJK}^{(-)} \quad (3.43)$$

This is all that is needed to split the theory into two parts. First I split the fundamental fields

$$E^{aI} = E^{(+aI)} + E^{(-aI)} \quad A_{aI} = A_{aI}^{(+)} + A_{aI}^{(-)} \quad (3.44)$$

Putting this into the Hamiltonian (3.36), yields

$$H^{tot} = \int_{\Sigma} d^3x \left(N^{(+)} \mathcal{H}^{(+)} + N^{(+a)} \mathcal{H}_a^{(+)} - A_{0I}^{(+)} \mathcal{G}^{(+I)} + c.c \right) \quad (3.45)$$

where

$$N^{(+)} := N + iN^* \quad (3.46)$$

$$N^{(+a)} := N^a + iN^{*a} \quad (3.47)$$

$$A_{0I}^{(+)} := \delta_I^{(+J)} A_{0J} \quad (3.48)$$

$$\mathcal{H}^{(+)} := \frac{1}{4} \epsilon_{abc} f_{IJK}^{(+)} E^{(+aI)} E^{(+bJ)} (B^{(+cK)} + \frac{2\Lambda i}{3} E^{(+cK)}) \approx 0 \quad (3.49)$$

$$\mathcal{H}_a^{(+)} := \frac{1}{2} \epsilon_{abc} E^{(+bI)} B_I^{(+c)} \approx 0 \quad (3.50)$$

$$\mathcal{G}_I^{(+)} := \mathcal{D}_a E^{(+aI)} = \partial_a E^{(+aI)} + f^{(+I)}_{JK} A_a^{(+J)} E^{(+aK)} \approx 0 \quad (3.51)$$

We also have

$$\begin{aligned} \{A_{aI}^{(+)}(x), E^{(+bJ)}(y)\} &= \delta_a^b \delta_I^{(+J)} \delta^3(x-y), \quad \{A_{aI}^{(-)}(x), E^{(-bJ)}(y)\} = \delta_a^b \delta_I^{(-J)} \delta^3(x-y), \\ \{A_{aI}^{(-)}(x), E^{(+bJ)}(y)\} &= \{A_{aI}^{(+)}(x), E^{(-bJ)}(y)\} = 0 \end{aligned} \quad (3.52)$$

And since all the fields in the $so(1,3)$ theory were taken real, the complex conjugate of the self dual part equals the antiself dual part. Thus, we see that the $so(1,3)$ theory really equals the complex Ashtekar formulation *without* the reality conditions. That is; complex Einstein gravity. If we had started with the algebras $so(4)$ or $so(2,2)$ instead of $so(1,3)$, we would have found two identical copies of the same real $so(3)$ or $so(1,2)$ theory. That is two copies of Einstein gravity for Euclidean or "ultra-hyperbolic" spacetime.

With this knowledge, that the $so(1,3)$ theory just is a real formulation of complex Einstein gravity, it is an easy task to answer all the questions that were asked in the end of the previous subsection: 1. The theory describes complex Einstein gravity. 2. The theory

is four-dimensionally diffeomorphism invariant. 3. The geometrical interpretation of the fields is that

$$\tilde{g}^{\alpha\beta} = \sqrt{-g}g^{\alpha\beta} = \begin{pmatrix} -\frac{1}{N^{(+)}} & \frac{N^{(+)}_a}{N^{(+)}} \\ \frac{N^{(+)}_a}{N^{(+)}} & -\frac{N^{(+)}}{8}E^{(+)}_{aI}E^{(+)}_I{}^b - \frac{N^{(+)}_aN^{(+)}_b}{N^{(+)}} \end{pmatrix} \quad (3.53)$$

is the densitized complex spacetime metric. This follows from the equivalence with the Ashtekar formulation, and formula for the spacetime metric in Ashtekar's variables. See [9]. 4. The symmetries generated by $\mathcal{H}_a^*$ and $\mathcal{H}^*$ is complex diffeomorphisms. Therefore, any two metrics that only differ by a complex coordinate transformation are to be considered physically equivalent. To see this, consider *e.g.* the transformations generated by $\tilde{\mathcal{H}}_a^*$ on $E^{(+)}_{aI}$.

How is then the Ashtekar formulation of conventional real Einstein gravity embedded into this larger $so(1,3)$ theory? We know that the reality conditions that are used to select real general relativity from Ashtekar's variables are [1]:

$$Im(E^{(+)}_{aI}E^{(+)}_I{}^b) = 0, \quad Im(E^{(+)}_{cM}f^{(+)}_{MIN}E^{(+)}_{(bI}\mathcal{D}_cE^{(+)}_{a)N}) = 0 \quad (3.54)$$

$$Im(N^{(+)}) = 0, \quad Im(N^{(+)}_a) = 0 \quad (3.55)$$

These conditions are easily translated back into the real $so(1,3)$ formulation where they become

$$E^{aI}E^{*b}_I = 0, \quad E^{*cM}f_{MIN}E^{(bI}\mathcal{D}_cE^{a)N} = 0, \quad N^* = 0, \quad N^{*a} = 0 \quad (3.56)$$

This means that any field configuration that satisfies (3.56) initially will continue to do so under the time evolution. Note, however, that there is nothing automatic about the Lorentzian signature of the metric, neither in the Ashtekar formulation nor in this $so(1,3)$ theory. The Lorentzian case is selected by requiring $Re(E^{(+)}_{aI}E^{(+)}_I{}^b) < 0$ or $E^{aI}E^b_I < 0$ (By this, I mean all eigenvalues are negative definite.)

Note also the similarity between the "reality conditions" (3.56) and the second class constraints found in the Hamiltonian formulation of the Hilbert-Palatini Lagrangian [1], [9]. The first two conditions in (3.56) equal exactly these second class constraints, while the two last conditions in (3.56) imply that $\mathcal{H}^*$ and $\mathcal{H}_a^*$ drop out from the total Hamiltonian.

3.4 The Lagrangians

Here, I will derive two Lagrangians whose Hamiltonian formulations equal the one given by (3.36). This is actually a rather trivial task once one knows that the $so(1,3)$ theory splits into the complex Ashtekar formulation plus its complex conjugate: the total Lagrangians will just be the complex Lagrangians that give the complex Ashtekar formulation, plus their complex conjugate. The first order Lagrangian is the selfdual Hilbert-Palatini Lagrangian [5], [6], [9], and the second order Lagrangian is the CDJ-Lagrangian [7], [8], [9]. However, I will also show what the manifestly real $so(1,3)$ Lagrangians look like, and give the definition of the spacetime metric in terms of the configuration space variables.

The first order Lagrangian leading to the complex Ashtekar formulation is

$$\mathcal{L} = \frac{1}{2} \epsilon^{\alpha\beta\gamma\delta} e_{\alpha A} e_{\beta B} R_{\gamma\delta}^{(+)\,AB} (A_\epsilon^{CD}) \quad (3.57)$$

where $e_{\alpha A}$ is the complex tetrad and $R_{\gamma\delta}^{(+)\,AB}$ is the self dual part of the Riemann tensor. I will assume that the full spin-connection A_α^{AB} is real, meaning that the self dual and antiself dual connection will be related via complex conjugation. It may seem strange to allow the tetrad to be complex while the $so(1, 3)$ connection remains real. The reason why this is possible is that the important connection in (3.57) really is the self dual $so(1, 3)$ connection which necessarily is complex. Furthermore, as is easily proven, a generic self dual field can always be written as the self dual part of a *real* field, without loss of generality.

To find the manifestly real formulation, I need to add the complex conjugate of the Lagrangian to the Lagrangian

$$\mathcal{L}^{tot} = \mathcal{L} + \bar{\mathcal{L}} = \frac{1}{2} \epsilon^{\alpha\beta\gamma\delta} e_{\alpha A} e_{\beta B} R_{\gamma\delta}^{(+)\,AB} (A_\epsilon^{CD}) + \frac{1}{2} \epsilon^{\alpha\beta\gamma\delta} \bar{e}_{\alpha A} \bar{e}_{\beta B} R_{\gamma\delta}^{(-)\,AB} (A_\epsilon^{CD}) \quad (3.58)$$

where $\bar{e}_\alpha^A$ is the complex conjugate of the tetrad. Now, I define the real objects

$$\begin{aligned} R_{\alpha\beta}^{AB} &= R_{\alpha\beta}^{(+)\,AB} + R_{\alpha\beta}^{(-)\,AB}, & R_{\alpha\beta}^{*\,AB} &= i \left(R_{\alpha\beta}^{(+)\,AB} - R_{\alpha\beta}^{(-)\,AB} \right) \\ V_\alpha^A &:= \frac{1}{2} (e_\alpha^A + \bar{e}_\alpha^A), & W_\alpha^A &:= \frac{-i}{2} (e_\alpha^A - \bar{e}_\alpha^A) \end{aligned} \quad (3.59)$$

With this, the Lagrangian becomes

$$\mathcal{L}^{tot} = \frac{1}{2} \epsilon^{\alpha\beta\gamma\delta} (V_{\alpha A} V_{\beta B} - W_{\alpha A} W_{\beta B}) R_{\gamma\delta}^{AB} (A_\epsilon^{CD}) + \epsilon^{\alpha\beta\gamma\delta} V_{\alpha A} W_{\beta B} R_{\gamma\delta}^{*\,AB} (A_\epsilon^{CD}) \quad (3.60)$$

which is the manifestly real $so(1, 3)$ Lagrangian for complex gravity. The complex metric is given by

$$g_{\alpha\beta} = e_{\alpha A} e_{\beta}^A = V_{\alpha A} V_{\beta}^A - W_{\alpha A} W_{\beta}^A + i V_{(\alpha A} W_{\beta)}^A \quad (3.61)$$

There is one strange feature of the Lagrangian (3.60). If one counts the number of components in $V_{\alpha A}$ and $W_{\alpha A}$ one gets 32, while the corresponding fields in the Hamiltonian formulation – N , N^* , N^a , N^{*a} and E^{aI} – only have 26 components. There are six components missing! A similar loss happens when one goes from the self dual Hilbert-Palatini Lagrangian to the Ashtekar formulation. In the Lagrangian, the tetrad $e_{\alpha A}$ has 16 components, but at the Hamiltonian level, we only find N , N^a and E^{ai} ($so(3)$) – 13 components. Three are missing! For that case, the three missing components are the ones that can be gauged away via antiself dual Lorentz transformations. That is, the tetrad transforms under the full Lorentz group, while the self dual antisymmetric products of two tetrads only transforms under the self dual part of the Lorentz Lie-algebra. Therefore, without restricting the Hamiltonian formulation, one may remove three components of the tetrad. Does there exist a similar local symmetry for the Lagrangian (3.60)? Yes, besides the normal Lorentz symmetry we also have local invariance under

$$\begin{aligned} \delta V_{\alpha A} &= \Lambda_A^{*\,B} W_{\alpha B}, & \delta W_{\alpha A} &= -\Lambda_A^{*\,B} V_{\alpha B} \\ \delta R_{\gamma\delta}^{AB} &= \Lambda^A_C R_{\gamma\delta}^{CB} - \Lambda^B_C R_{\gamma\delta}^{CA} \end{aligned} \quad (3.62)$$

where $\Lambda^A{}_B$ belongs to $so(1, 3)$, and the $*$ denotes the dual of the field: $\Lambda_{AB}^* := \frac{1}{2}\epsilon_{AB}{}^{CD}\Lambda_{CD}$. Thus, we have a six-dimensional local symmetry for the Lagrangian (3.60) which is *not* represented by a first class constraint at the Hamiltonian level. This ends the discussion about the first order Lagrangian.

Finding a second order Lagrangian corresponding to a given Hamiltonian, means performing a Legendre transform. That is solve the momentum's equation of motion to get a solution for the momentum in terms of the phase space coordinates and their velocities. When there are constraints present, one also has the choice of treating the constraints as primary or secondary w.r.t the wanted Lagrangian. If the constraints are taken to be primary, the solution for the momenta should automatically satisfy the constraints, while in the case of secondary constraints no such algebraic features are required. In Legendre transforms from the Ashtekar formulation it has been found convenient to treat the vector constraint as primary while the Hamiltonian and the Gauss law constraint are considered to be secondary. For more details regarding this, see *e.g.* [8]. Thus, we use the (3.45) version of the $so(1, 3)$ Hamiltonian and calculate the equations of motion for $E^{(+a)I}$:

$$F_{0a}^{(+I)} = \frac{N^{(+)}}{2}\epsilon_{abc}E^{(+b)J}B^{(+c)K}f^{(+I)}{}_{JK} + \frac{1}{2}N^{(+b)}\epsilon_{bac}B^{(+c)I} \quad (3.63)$$

$$\mathcal{H}_a^{(+)} = \frac{1}{2}\epsilon_{abc}E^{(+b)I}B_I^{(+c)} = 0 \quad (3.64)$$

The equations for the antiself dual fields are exact copies of (3.63) and (3.64). To solve these equations for $E^{(+a)I}$, I will assume that the magnetic field is non-degenerate, or $\det(B^{(+a)I}B_I^{(+b)}) \neq 0$. If this is true, I can always expand any vector field in terms of $B^{(+a)I}$. For instance

$$E^{(+a)I} = \Psi^{IJ}B_J^{(+a)}. \quad (3.65)$$

Note that Ψ^{IJ} is self dual in both indices implying that the maximal rank of it is three. Using this expansion in (3.64) shows that Ψ^{IJ} must be symmetric. This, together with the formula $f_{IJK}^{(+)}B^{(+a)I}B^{(+b)J}B^{(+c)K} = |B^{(+d)I}| \epsilon^{abc}$, where $|B^{(+d)I}| := \frac{1}{6}f_{IJK}^{(+)}B^{(+a)I}B^{(+b)J}B^{(+c)K}\epsilon_{abc}$, makes it possible to rewrite (3.63) as

$$\Omega^{(+MI)} = \frac{1}{\eta^{(+)}} \left(\Psi^{MI} - g^{(+MI)}Tr(\Psi) \right) \quad (3.66)$$

where I have defined $\eta^{(+)} := -(2N^{(+)} |B^{(+d)I}|)^{-1}$ and $\Omega^{(+MI)} := 2B^{(+a)M}F_{0a}^{(+I)} = \epsilon^{\alpha\beta\delta\epsilon}F_{\alpha\beta}^{(+M)}F_{\delta\epsilon}^{(+I)}$. Eq. (3.66) is really just six of the nine components of eq. (3.63). The remaining three components can be used to fix $N^{(+a)}$, but that does not make any difference in the Legendre transform since the choice of treating $\mathcal{H}_a$ and $\mathcal{H}_a^*$ as primary means that N^a and N^{*a} completely drop out from the Lagrangian. Now, (3.66) is easily solved for Ψ^{IJ} , and the solution for $E^{(+a)I}$ becomes

$$E^{(+a)I} = \eta^{(+)} \left(\Omega^{(+IJ)} - \frac{1}{2}Tr(\Omega^{(+)}) \right) B_J^{(+a)} \quad (3.67)$$

The Lagrangian is $\mathcal{L} = E^{(+a)I}\dot{A}_{aI}^{(+)} + E^{(-a)I}\dot{A}_{aI}^{(-)} - H^{tot}$, and using (3.67) for $E^{(+a)I}$ and the complex conjugate of (3.67) for $E^{(-a)I}$, I get

$$\mathcal{L} = \frac{\eta^{(+)}}{8} \left(\Omega^{(+IJ)}\Omega_{IJ}^{(+)} - \frac{1}{2}(\Omega_I^{(+I)})^2 \right) + c.c \quad (3.68)$$

where as before, the complex conjugate of a self dual field equals the antiself dual field.

To find the real $so(1,3)$ Lagrangian from here, I need some formulas relating $\Omega^{(+IJ)}$ and $\Omega^{(-IJ)}$ to $\Omega^{IJ} := \epsilon^{\alpha\beta\delta\epsilon} F_{\alpha\beta}^I F_{\delta\epsilon}^J$ and $\Omega^{*IJ} := \Omega^{IK} g_K^*{}^J$, etc. By definition

$$\Omega^{(+IJ)} = \Omega^{KL} \delta_K^{(+I)} \delta_L^{(+J)} = \frac{1}{4} \left(\Omega^{IJ} - i^* \Omega^{IJ} - i \Omega^{*IJ} - * \Omega^{*IJ} \right). \quad (3.69)$$

Using this in the Lagrangian (3.68), yields

$$\begin{aligned} \mathcal{L} = & \frac{\eta}{8} \left(Tr(\Omega^2) - Tr(\Omega^* \Omega^*) - \frac{1}{2} (Tr \Omega)^2 + \frac{1}{2} (Tr \Omega^*)^2 \right) + \\ & \frac{\eta^*}{4} \left(Tr(\Omega \Omega^*) - \frac{1}{2} Tr \Omega Tr \Omega^* \right) \end{aligned} \quad (3.70)$$

where

$$\begin{aligned} \eta &:= \frac{1}{2} Re(\eta^{(+)}) = \frac{1}{4} (\eta^{(+)} + \eta^{(-)}) = -\frac{1}{2} \frac{|B^{aI}| N + |B^{*aI}| N^*}{(|B^{aI}|^2 + |B^{*aI}|^2)(N^2 + (N^*)^2)} \\ \eta^* &:= \frac{1}{2} Im(\eta^{(+)}) = \frac{1}{4i} (\eta^{(+)} - \eta^{(-)}) = -\frac{1}{2} \frac{|B^{*aI}| N - |B^{aI}| N^*}{(|B^{aI}|^2 + |B^{*aI}|^2)(N^2 + (N^*)^2)} \\ Tr(\Omega^* \Omega^*) &:= \Omega^{IJ} \Omega^{KL} g_{IK}^* g_{JL}^*, \quad etc \end{aligned} \quad (3.71)$$

This completes the Legendre transform. The Lagrangian density (3.70) is the Lagrangian whose Hamiltonian is given by (3.36). Furthermore, from the explicit form of the Lagrangian, it is obvious that the theory is diffeomorphism invariant.

When it comes to identifying a spacetime metric in this theory, we can again rely on the results from the conventional Ashtekar/CDJ-formulation. See for instance [7], [8]. The spacetime metric is

$$\tilde{g}^{\alpha\beta} = \sqrt{-g} g^{\alpha\beta} = \frac{2\eta^{(+)}}{3} K^{(+)\alpha\beta} \quad (3.72)$$

where

$$K^{(+)\alpha\beta} := \epsilon^{\alpha\gamma\delta\epsilon} \epsilon^{\beta\kappa\tau\rho} F_{\gamma\delta}^I F_{\epsilon\rho}^J F_{\kappa\tau}^K f_{IJK}^{(+)} \quad (3.73)$$

4 The loop-algebra theory

In the search for a Lie-algebra that provides us with the structure constant identity (2.5), one is naturally led to extensions of the finite dimensional Lie-algebras ($so(3)$, $so(1,3)$, etc.) that are known to satisfy the identity. The loop-algebras are infinite dimensional examples of such extensions.

In this section, I will study an Ashtekar-like canonical theory based on the loop-algebra $\widetilde{so(3)}$. The construction can easily be generalized to the algebras $so(1,2)$, $so(4)$, $so(1,3)$, as well as isomorphic ones. In subsection 4.1, I give a short description of the algebra. Subsection 4.2 gives the Hamiltonian description, and, finally, in subsection 4.3 I show that the theory can be seen as an infinite number of copies of the Ashtekar formulation.

4.1 The loop-algebra

For details regarding loop-algebras and Kac-Moody algebras, see *e.g* [10].

Given a finite dimensional Lie-algebra $\mathcal{A}$, there exist an infinite dimensional extension of it $\tilde{\mathcal{A}}$. If T_I are the generators of the finite dimensional base-algebra, they satisfy the relation $[T_I, T_J] = f_{IJ}^K T_K$. In the extended algebra, we have the generators T_I^n , where $T_I^0 = T_I$ and the new index n denote the level of the generator, it takes all integer values. In the extended algebra, the Lie-bracket is

$$[T_I^n, T_J^m] = f^{nm}_k f_{IJ}^K T_K^k = f_{IJ}^K T_K^{n+m} \quad (4.1)$$

where we see that the structure constant f^{nm}_k is non-vanishing only for $n + m = k$. One may notice that it is not possible to stop at any finite level, except the zeroth level, since the Lie-bracket produces higher levels. However, one is allowed to use only the even levels since the sum of two even integers is even. Often, in the context of loop-algebras, one consider the central extension of the algebra. In that case, the right-hand side of (4.1) gets an additional term $n\delta^{n,-m} g_{IJ} c$, where c is the one dimensional center – it commutes with all elements – and g_{IJ} is a bilinear form of the base Lie-algebra. Furthermore, it is straightforward to check that the centrally extended loop-algebra also allows an extension with an element d :

$$[d, T_I^n] = nT_I^n, \quad [d, d] = 0, \quad [c, d] = 0 \quad (4.2)$$

The resulting Lie-algebra $\hat{\mathcal{A}} = \tilde{\mathcal{A}} \oplus \mathcal{R}c \oplus \mathcal{R}d$ is called an affine Kac-Moody algebra.

Now, the name loop-algebra comes from the fact that the algebra admits a very nice interpretation; it can be seen as $\mathcal{A}$ -algebra valued fields on the circle, S^1 . That is, given a $\mathcal{A}$ -algebra valued field $h(\theta)$ that is a function of a coordinate θ on S^1 we may make a fourier series expansion of it ($h(\theta + 2\pi) = h(\theta)$)²:

$$h(\theta) = \sum_{n=-\infty}^{\infty} h_n \exp(-in\theta), \quad h_n := \frac{1}{2\pi} \int_0^{2\pi} d\theta h(\theta) \exp(in\theta) \quad (4.3)$$

Thus, we see that we may write $h(\theta) = h_n^I T_I^n := h_n^I T_I \exp(-in\theta)$, and from there, the commutation relation (4.1) follows immediately. We also see that d can be represented as $i \frac{d}{d\theta}$. This way of representing $\tilde{\mathcal{A}}$ will be used in subsection 4.3 to get a better understanding of the theory. Note, however, that we are not forced to use this representation, we may just see the Lie-algebra $\tilde{\mathcal{A}}$ as abstractly defined by (4.1).

Now, we need an invariant bilinear form. Here, I will only study the loop-algebra $\tilde{\mathcal{A}}$, and at the end comment on the case were one extends it with c and d . As in the previous section, the form is invariant if

$$\langle [C, A], B \rangle + \langle A, [C, B] \rangle = 0, \quad (4.4)$$

where $\langle \cdot, \cdot \rangle$ and $[\cdot, \cdot]$ denote the bilinear form and the Lie-bracket, respectively. A , B , and C belongs to $\tilde{\mathcal{A}}$ Translating this condition to the structure constants, it becomes:

$$f^{nm}_k f_{IJ}^K \tilde{g}_{KL} + f^{nl}_k f_{IL}^K \tilde{g}_{KJ}^{km} = 0 \quad (4.5)$$

²Sometimes one considers a slightly weaker periodicity condition $h(\theta + 2\pi) = \tau(h(\theta))$, where τ is an (outer) automorphism of the Lie-algebra $\mathcal{A}$. The resulting algebra is called a twisted loop-algebra/Kac-Moody algebra

where I have defined $\tilde{g}_{IJ}^{nm} := \langle T_I^n, T_J^m \rangle$. Using that $f^{nm}_k = \delta(n+m-k)$ eq. (4.5) gives that the bilinear form must be of the form $\tilde{g}_{IJ}^{nm} = \tilde{g}^{nm} g_{IJ}$ where g_{IJ} is an invariant bilinear form for $\mathcal{A}$, the invariance condition becomes:

$$f^{nmk} - f^{nkm} = 0 \quad (4.6)$$

Here, I have raised the "level-index" with $\tilde{g}^{nm}$. Since we know that f^{nm}_k has the value one if $n+m=k$, and is zero otherwise, we can easily transfer the condition (4.6) to $\tilde{g}^{nm}$:

$$\tilde{g}^{n,m} = \tilde{g}^{n+k,m-k} \quad (4.7)$$

If one sees $\tilde{g}^{nm}$ as an infinite dimensional matrix, the equation (4.7) says that all elements on the same "cross-diagonal" must be equal. The standard choice for this bilinear form is $\tilde{g}^{nm} = \delta(n+m)$, which satisfy the required relation (4.7). Furthermore, if we introduce the extensions c and d , this choice of bilinear form is unique if we require invariance under conjugation with d . I will, however, not make this choice here, instead I notice that there exist a basis for all invariant bilinear forms. That is, every invariant bilinear form $\tilde{g}^{nm}$ may be written as

$$\tilde{g}^{nm} = \sum_{k=-\infty}^{\infty} a_{(k)} \tilde{g}_{(k)}^{nm} \quad (4.8)$$

$$\tilde{g}_{(k)}^{nm} = \delta(-k+n+m) \quad (4.9)$$

where the $a_{(k)}$ are taken to be real numbers. Notice the similarity between f^{nm}_k and $\tilde{g}_{(k)}^{nm}$. Thus, every bilinear invariant form of the Lie-algebra may be written as

$$\tilde{g}^{nm} = v^k f^{nm}_k \quad (4.10)$$

for some Lie-algebra valued field v^k . To be able to raise and lower the "level-indices" n , m , ..., I pick out one bilinear form and its inverse to be responsible for that operation: $\tilde{g}_{(0)}^{nm} = f^{nm}_0$ and $\tilde{g}_{(0)nm} \tilde{g}_{(0)}^{mk} = \delta_n^k E.g.$

$$f^{nmk} := f^{nm} \tilde{g}_{(0)}^{lk}, \quad \tilde{g}_{(k)n}^m := \tilde{g}_{(k)}^{km} \tilde{g}_{(0)nk} \quad (4.11)$$

The following relations will then hold

$$\tilde{g}_{(k)}^{nm} = \delta(-k+n+m), \quad \tilde{g}_{(k)n}^m = \delta(k+n-m), \quad \tilde{g}_{(k)nm} = \delta(k+n+m) \quad (4.12)$$

Translating all these bilinear forms to the S^1 -representation, we see that the bilinear form $\tilde{g}_{(k)}^{nm}$ corresponds to integrating over the circle with the weight-factor $\exp(ik\theta)$. That is, if $h(\theta)$ and $f(\theta)$ belongs to $\tilde{\mathcal{A}}$, we get

$$\langle h(\theta), f(\theta) \rangle_{(k)} = h_n^I f_m^J \langle T_I^n, T_J^m \rangle_{(k)} = h_n^I f_m^J \tilde{g}_{(k)}^{nm} g_{IJ} = \sum_{n=-\infty}^{\infty} h_{Jn} f_{k-n}^J \quad (4.13)$$

which corresponds to

$$\langle h(\theta), f(\theta) \rangle_{(k)} = \frac{1}{2\pi} \int_0^{2\pi} d\theta \text{Tr}(h(\theta) f(\theta)) \exp(ik\theta) \quad (4.14)$$

where I have written the bilinear form in $\mathcal{A}$ as a trace: $\langle T_I, T_J \rangle := \text{Tr}(T_I T_J)$. From (4.14) it is also clear that the bilinear form is not invariant under conjugation with d :

$$\begin{aligned} \delta^d \langle h(\theta), f(\theta) \rangle_{(k)} &= \langle [d, h(\theta)], f(\theta) \rangle + \langle h(\theta), [d, f(\theta)] \rangle = i \langle h'(\theta), f(\theta) \rangle + \\ & i \langle h(\theta), f'(\theta) \rangle = \frac{i}{2\pi} \int_0^{2\pi} d\theta \text{Tr} (h'(\theta) f(\theta) + h(\theta) f'(\theta)) \exp(ik\theta) = \\ & k \langle h(\theta), f(\theta) \rangle_{(k)} \end{aligned} \quad (4.15)$$

Thus, we see that the bilinear form is only invariant under d -transformations iff $k = 0$. For $k \neq 0$, we instead get something like a "covariant bilinear form".

Finally, before I start constructing the theory, I need to show that this algebra provides a structure constant identity of the type (2.5). To do this, I first pick a base-algebra $\mathcal{A}$ that by itself has structure constants that satisfy this type of identity. The simplest example is $\mathcal{A} = \mathfrak{so}(3)$, where $f_{IJ}{}^K f_{KLM} = \delta_{IL} \delta_{JM} - \delta_{JL} \delta_{IM}$ with a proper choice of basis. For $\widetilde{\mathfrak{so}(3)}$, we get

$$f_{IJ}{}^K f_{KLM} f^{nm}{}_k f^{klj} = (\delta_{IL} \delta_{JM} - \delta_{JL} \delta_{IM}) \sum_{k=-\infty}^{\infty} \tilde{g}_{(k)}^{nm} \tilde{g}_{(-k)}^{lj} \quad (4.16)$$

which is an identity of the requested type. Furthermore, from (4.12) it is clear that $\tilde{g}_{(k)n}^j \tilde{g}_{(m)}^{nl} = \tilde{g}_{(k+m)}^{jl}$, which is the second requirement that is needed; the product of two bilinear forms must be an invariant bilinear form.

4.2 The $\widetilde{\mathfrak{so}(3)}$ Hamiltonian

In this subsection, I will construct a gauge and diffeomorphism invariant theory based on the Lie-algebra $\widetilde{\mathfrak{so}(3)}$. The construction will be very similar to the $\mathfrak{so}(1,3)$ theory presented in subsection 3.2. The major difference is that we now have an infinite number of invariant bilinear forms for $\widetilde{\mathfrak{so}(3)}$ compared to only two for $\mathfrak{so}(1,3)$. This will imply the introduction of an infinite number of new constraints. I will only treat the non-extended $\widetilde{\mathfrak{so}(3)}$ here, and the reason why I do not include the central extension and the d -generator is that I have not been able to find a closed constraint algebra for that case.

The basic fields are an $\widetilde{\mathfrak{so}(3)}$ connection A_{aI}^n and its conjugate momentum, E_n^{aI} . They satisfy

$$\{A_{aI}^n(x), E_m^{bJ}(y)\} = \delta_a^b \delta_I^J \delta_m^n \delta^3(x - y) \quad (4.17)$$

To write out the constraints, I first define: $E_{(k)n}^{aI} := \tilde{g}_{(k)n}^m E_m^{aI}$, and analogous for $A_{(k)a}^{In}$. (If one sees E_n^{aI} as an infinite dimensional vector of three-by-three matrices, $E_{(k)n}^{aI}$ is just E_n^{aI} shifted k steps along the vector.) Here are the constraints

$$\mathcal{G}^{In} := \mathcal{D}_a E^{aIn} = \partial_a E^{aIn} + f^I{}_{JK} f^{nmk} A_{am}^J E_k^{aK} \approx 0 \quad (4.18)$$

$$\mathcal{H}_{(k)a} = \frac{1}{2} \epsilon_{abc} E_{(k)I}^{bn} E_n^{cI} \approx 0 \quad (4.19)$$

$$\mathcal{H}_{(k)} = \frac{1}{4} \epsilon_{abc} f_I{}^{JK} f^n{}_{mk} E_{(k)J}^{bm} E_K^{ck} \left(B_n^{aI} + \frac{2\Lambda}{3} E_n^{aI} \right) \approx 0 \quad (4.20)$$

$$(4.21)$$

where $B_n^{aI} := \epsilon^{abc} F_{bcn}^I = \epsilon^{abc} (2\partial_b A_{cn}^I + f_{JK}^I f^{mk}{}_n A_{bm}^J A_{ck}^K)$. Now it is straightforward to go through the poisson bracket calculations in the same manner as in section 3.2: **I.** Check that $\mathcal{G}_I^n$ and $\tilde{\mathcal{H}}_{(k)a} := \mathcal{H}_{(k)a} - A_{(k)a}^I \mathcal{G}_I^n$ generate gauge transformations and spatial diffeomorphisms times a k -shift, respectively. **II.** Calculate all poisson brackets containing these two constraints by using the transformation properties of the constraints. **III.** Calculate the remaining poisson brackets $\{\mathcal{H}_{(k)}[N], \mathcal{H}_{(m)}[M]\}$.

I. The transformations generated by $\mathcal{G}^{In}$ and $\tilde{\mathcal{H}}_{(k)a} := \mathcal{H}_{(k)a} - A_{(k)a}^{In} \mathcal{G}_{In}$ are

$$\begin{aligned} \delta^{\mathcal{G}^{In}} E_m^{aI} &:= \{E_m^{aI}, \mathcal{G}_n^J[\Lambda_J^n]\} = f^I{}_{JK} f_{mn}{}^l \Lambda^{Jn} E_l^{aK} \\ \delta^{\mathcal{G}^{In}} A_{am}^I &:= \{A_{am}^I, \mathcal{G}_n^J[\Lambda_J^n]\} = -\mathcal{D}_a \Lambda_m^I \\ \delta^{\tilde{\mathcal{H}}_{(k)a}} E_n^{aI} &:= \{E_n^{aI}, \tilde{\mathcal{H}}_{(k)b}[N^b]\} = N^b \partial_b E_{(k)n}^{aI} - E_{(k)n}^{bI} \partial_b N^a + E_{(k)n}^{aI} \partial_b N^b = \mathcal{L}_{N^b} E_{(k)n}^{aI} \\ \delta^{\tilde{\mathcal{H}}_{(k)a}} A_a^{In} &:= \{A_a^{In}, \tilde{\mathcal{H}}_{(k)b}[N^b]\} = N^b \partial_b A_{(k)a}^{In} + A_{(k)b}^{In} \partial_a N^b = \mathcal{L}_{N^b} A_{(k)a}^{In} \end{aligned} \quad (4.22)$$

which shows that $\mathcal{G}^{In}$ generates gauge transformations and $\tilde{\mathcal{H}}_{(k)a}$ generates spatial diffeomorphisms times a k -shift.

II. With this knowledge, the calculation of all Poisson brackets containing $\mathcal{G}_n^I$ and $\tilde{\mathcal{H}}_a^{(k)}$ simplifies significantly. We know that $\mathcal{G}_n^I, \mathcal{H}_{(k)}$ are gauge-covariant, and that all constraints are diffeomorphism covariant. This gives

$$\{\mathcal{G}_n^I[\Lambda_I^n], \mathcal{G}^{Jm}[\Gamma_J^m]\} = \mathcal{G}_n^K[f_{KIJ} f^{knm} \Lambda_n^I \Gamma_m^J] \quad (4.23)$$

$$\{\mathcal{G}_n^I[\Lambda_I^n], \tilde{\mathcal{H}}_{(k)a}[N^a]\} = \mathcal{G}_n^I[-\mathcal{L}_{N^a} \Lambda_{(k)I}^n] \quad (4.24)$$

$$\{\mathcal{G}_n^I[\Lambda_I^n], \mathcal{H}^{(k)}[N]\} = 0 \quad (4.25)$$

$$\{\tilde{\mathcal{H}}_{(k)a}[N^a], \tilde{\mathcal{H}}_{(l)b}[M^b]\} = \tilde{\mathcal{H}}_{(k+l)a}[-\mathcal{L}_{M^b} N^a] \quad (4.26)$$

$$\{\mathcal{H}_{(k)}[N], \tilde{\mathcal{H}}_{(l)a}[N^a]\} = \mathcal{H}_{(k+l)}[-\mathcal{L}_{N^a} N] \quad (4.27)$$

III. To calculate $\{\mathcal{H}_{(k)}[N], \mathcal{H}_{(m)}[M]\}$, the identity (4.16) comes in handy. The result is

$$\{\mathcal{H}_{(m)}[N], \mathcal{H}_{(n)}[M]\} = \sum_{k=-\infty}^{\infty} \mathcal{H}_{(k+m+n)a} [q_{(-k)}^{ab} (N \partial_b M - M \partial_b N)] \quad (4.28)$$

where I have defined $q_{(k)}^{ab} := -E_{(k)n}^{aI} E_I^{bn}$. This completes the constraint analysis. All constraint are first class, and the total Hamiltonian is

$$H^{tot} = \int_{\Sigma} d^3x \left(-A_{0I}^n \mathcal{G}_n^I + \sum_{k=-\infty}^{\infty} (N^{(k)} \mathcal{H}_{(k)} + N^{(k)a} \mathcal{H}_{(k)a}) \right) \quad (4.29)$$

where $A_{0I}^n, N_{(k)}$ and $N_{(k)}^a$ are Lagrange multipliers. Thus, we see that it is indeed possible to construct a consistent Hamiltonian formulation based on the Lie-algebra $\widetilde{so(3)}$. However, in doing so, one is forced to include an infinite number of constraints. The question is now; have we gained any local degrees of freedom by enlarging the gauge algebra from $so(3)$ to $\widetilde{so(3)}$? This is not a trivial question to answer in this description, since we have both an infinite number of phase space variables as well as an infinite number of constraints. If we naively try to calculate the number of local degrees of freedom by truncating the theory at some finite level N , we seem to get more constraints than phase

space variables, meaning that we have no local degrees of freedom. However, we know that that is not true, we know that the non-trivial $so(3)$ Ashtekar formulation of Einstein gravity is contained in this theory. So, for some reason the naive count does not give the complete answer. One reason could be that by truncating the theory at a finite level, we actually break the gauge invariance of the theory, meaning that the Gauss law constraint cannot any longer be treated as a first class constraint. (The same will also be true for some of the other constraints.) However, as I will show in the next subsection, by rewriting this theory as a theory with a finite number of constraints on every point on the circle, it is clear that this theory actually has an infinite number of degrees of freedom per point in spacetime. So, what have we gained by enlarging the conventional Ashtekar formulation in this way? My starting motivation for enlarging the theory in the first place was the hope of finding a real theory for Lorentzian spacetime. Can this be accomplished with the theory presented here? To answer that question, one really needs to know how the metric is defined in terms of the fields in this formulation. Remember that the need of complex fields in the conventional $so(3)$ Ashtekar formulation, can be seen as coming from the requirement that $E^{aI} E_I^b$ should be negative definite [4]. And, as is shown in the next subsection, this $\widetilde{so(3)}$ theory is really equivalent to an infinite number of copies of complex Einstein gravity, meaning that the theory actually has an infinite number of complex spacetime metrics. Also, at this stage it is not even clear that the conventional theory of real Einstein gravity is contained in the *real* $\widetilde{so(3)}$ formulation. It is obvious that it is contained in the *complex* $\widetilde{so(3)}$ theory; just put all higher modes in the fields to zero, and we recover the conventional complex Ashtekar formulation. It is, however, true that conventional real Einstein gravity is contained also in the *real* $\widetilde{so(3)}$ theory, and to prove that, I will show how we can recover the ADM-Hamiltonian from the $\widetilde{so(3)}$ Hamiltonian.

Consider a field configuration in the $\widetilde{so(3)}$ theory that is initially of the form:

$$E_n^{aI} = E^{aI} g_n, \quad g_n := \frac{1}{\pi n}((-1)^n - 1) \quad (4.30)$$

$$A_{aIn} = \delta_n^0 \Gamma_{aI}(\epsilon^{bJ}) - K_{aI} g_n \quad (4.31)$$

$$N_{(k)} = N \delta_k^0 \quad (4.32)$$

$$N_{(k)} = N^a \delta_k^0 \quad (4.33)$$

$$A_{0K}^n = -\Lambda_K \delta_0^n + L_K g^n \quad (4.34)$$

where Γ_{aI} is the spin-connection defined to annihilate E^{aI} :

$$D_a E^{bI} = \partial_a E^{bI} + \Gamma_{ac}^b E^{cI} + f^{IJK} \Gamma_{aJ} E^{bK} - \Gamma_{ac}^c E^{bI} = 0 \quad (4.35)$$

Here I have also introduced the E^{ai} compatible Christoffel-connection Γ_{ab}^c . Both Γ_{ab}^c and Γ_{ai}^c can be uniquely solved for from (4.35) above. See for instance [9]. Note that g_n are the Fourier coefficients of the imaginary step function:

$$\sum_{n=-\infty}^{\infty} g_n \exp(-in\theta) = \begin{cases} i, & 0 < \theta < \pi \\ -i, & \pi < \theta < 2\pi \end{cases} \quad (4.36)$$

Now, to check that this field configuration will not change under time evolution, and also to rewrite the constraints, we will need the identity

$$f^{nmk} g_m g_k = -1 \delta_0^n \quad (4.37)$$

From (4.37) it follows that $g_n g^n = -1$, $f^{nmk} g_n g_m g_k = 0$ and $f^{nmk} g_m g_k f_n^{jl} g_j g_l = 1$. Writing down the equations of motion for E_n^{aI} and A_{aI}^n we get from the Hamiltonian (4.29), and using the field configurations (4.30)-(4.34), yields

$$\begin{aligned} \dot{E}_n^{aI} = & \dots = -\delta_n^0 f^{IJK} E_J^b E_K^a D_b N + g_n N K_{bN} E^{b[I} E^{aN]} + g_n E^{[aI} D_b N^{b]} + \\ & \delta_n^0 f^{IJK} K_{bJ} N^{[b} E_K^{a]} - g_n \Lambda_K f^{KIJ} E_J^a - \delta_n^0 L_K f^{KIJ} E_J^a \end{aligned} \quad (4.38)$$

$$\begin{aligned} \dot{A}_{aI}^n = & \dots = N f_{IJK} E^{bJ} \left(R_{ab}^K(\mathbf{r}) g^n + D_{[a} K_{b]}^K \delta_0^n - f^{KLM} K_{aL} K_{bM} g^n \right) + \\ & N^b \left(R_{baI}(\mathbf{r}) \delta_0^n - D_{[b} K_{a]I} g^n - f_{IJK} K_b^J K_a^K \delta_0^n \right) + g^n D_a L_I - \\ & \delta_0^n f_{IJK} \Gamma_a^J \Lambda^K + g^n f_{IJK} K_a^J \Lambda^K + \delta_0^n f_{IJK} K_a^J L^K \end{aligned} \quad (4.39)$$

where $R_{ab}^I := \partial_{[a} \Gamma_{b]}^I + f^{IJK} \Gamma_{aJ} \Gamma_{bK}$. From (4.38) it is clear that the parts that seem to break the initial field configuration (4.30) are just gauge transformations plus a term that vanishes due to the Gauss law constraint (4.42). However, to simplify things further, we may fix this freedom also, by choosing:

$$L^I = E^{bI} D_b N + N^b K_b^I \quad (4.40)$$

With this choice, E_n^{aI} will maintain its form under the time evolution. When it comes to A_{aI}^n , we must require that its zero-mode continue to annihilate E^{aI} during the time evolution. That is

$$\{D_a E^{bI}, H^{tot}\} = D_a \dot{E}^{bI} + \dot{\Gamma}_{ac}^b E^{cI} - \dot{\Gamma}_{ac}^c E^{bI} + f^{IJK} \delta_0^n \dot{A}_{aJn} E_K^b = 0 \quad (4.41)$$

It is straightforward, although rather tedious, to put (4.38) and (4.39) into (4.41) and show that the initial field configuration is indeed maintained under the time evolution. Furthermore, using this field configuration in the constraints, we get

$$\mathcal{G}^{In} = \delta_0^n f^{IJK} K_{aJ} E_K^a \quad (4.42)$$

$$\begin{aligned} \mathcal{H}_{(k)a} = & g_{(k)} E^{bI} R_{abI}(\mathbf{r}_{cK}) + \delta_{(k)}^0 E^{bI} D_{[a} K_{b]I} - g_{(k)} E_I^b f^{IJK} K_{aJ} K_{bK} \approx \\ & \delta_{(k)}^0 E^{bI} D_{[a} K_{b]I} \end{aligned} \quad (4.43)$$

$$\begin{aligned} \mathcal{H}_{(k)} = & -\delta_{(k)}^0 \frac{1}{2} f_{IJK} E^{aI} E^{bJ} (R_{ab}^K(\mathbf{r}_{cK}) - f^{KLM} K_{aL} K_{bM}) + g_{(k)} f_{IJK} E^{aI} E^{bJ} D_a K_b^K + \\ & \approx \frac{1}{2} \delta_{(k)}^0 \left(-f_{IJK} E^{aI} E^{bJ} R_{ab}^K(\mathbf{r}_{cK}) + E^{aI} E^{bJ} K_{a[I} K_{bJ]} \right) \end{aligned} \quad (4.44)$$

where I have used the Bianchi identity and the Gauss law in (4.43), and the Gauss law in (4.44). But, these constraints are just the conventional ADM-constraints for triad gravity. This means that every solution to Einstein's equations is also a solution to the real $so(3)$ theory. However, the price we have to pay here to get a real Ashtekar-like theory, containing Einstein gravity, is rather high; we had to introduce an infinite number of constraints as well as phase space variables, per point. Also, if the example above shows a generic feature of the inclusion of Lorentzian gravity into the real $so(3)$ theory, we must allow the fields to have "infinitely long tails" – all odd higher modes have to be non-zero – and with a very slow convergence. Of course, the field configurations chosen here, to show the relation to GR, is only one example, and it may be possible to find nicer field configurations that do the job. In fact, as I will show in the next subsection, the above mentioned problems with this theory can be solved by considering a theory based on a finite dimensional version of the Lie-algebra $so(3)$.

4.3 The $\widetilde{so(3)}$ theory on a circle

As was shown in subsection 4.1, one may see a general element in the loop-algebra $\widetilde{so(3)}$ as an $so(3)$ valued function on the circle. That is

$$E^{aI}(\theta) := \sum_{n=-\infty}^{\infty} E_n^{aI} \exp(-in\theta), \quad \Rightarrow \quad E_n^{aI} = \frac{1}{2\pi} \int_0^{2\pi} d\theta E^{aI}(\theta) \exp(in\theta) \quad (4.45)$$

$$A_a^I(\theta) := \sum_{n=-\infty}^{\infty} A_{an}^I \exp(-in\theta), \quad \Rightarrow \quad A_{an}^I = \frac{1}{2\pi} \int_0^{2\pi} d\theta A_a^I(\theta) \exp(in\theta) \quad (4.46)$$

$$N(\theta) := \sum_{k=-\infty}^{\infty} N_{(k)} \exp(-ik\theta), \quad \Rightarrow \quad N_{(k)} = \frac{1}{2\pi} \int_0^{2\pi} d\theta N(\theta) \exp(ik\theta) \quad (4.47)$$

$$N^a(\theta) := \sum_{k=-\infty}^{\infty} N_{(k)}^a \exp(-ik\theta), \quad \Rightarrow \quad N_{(k)}^a = \frac{1}{2\pi} \int_0^{2\pi} d\theta N^a(\theta) \exp(ik\theta) \quad (4.48)$$

$$A_0^I(\theta) := \sum_{n=-\infty}^{\infty} A_{0n}^I \exp(-in\theta), \quad \Rightarrow \quad A_{0n}^I = \frac{1}{2\pi} \int_0^{2\pi} d\theta A_0^I(\theta) \exp(in\theta) \quad (4.49)$$

Using this in the fundamental Poisson bracket, we get

$$\{A_{aI}(x, \theta), E^{bJ}(y, \theta')\} = \delta_a^b \delta_I^J \delta^3(x - y) \sum_{n=-\infty}^{\infty} \exp(-in(\theta - \theta')) = 2\pi \delta_a^b \delta_I^J \delta^3(x - y) \delta(\theta - \theta') \quad (4.50)$$

and the total Hamiltonian becomes

$$H^{tot} = \frac{1}{2\pi} \int_{\Sigma} d^3x \int_0^{2\pi} d\theta \left(N(\theta) \mathcal{H}(\theta) + N^a(\theta) \mathcal{H}_a(\theta) - A_{0I}(\theta) \mathcal{G}^I(\theta) \right) \quad (4.51)$$

where

$$\mathcal{H}(\theta) = \frac{1}{4} \epsilon_{abc} f_{IJK} E^{aI}(\theta) E^{bJ}(\theta) B^{cK}(\theta) \quad (4.52)$$

$$\mathcal{H}_a(\theta) = \frac{1}{2} \epsilon_{abc} E^{bI}(\theta) B_I^c(\theta) \quad (4.53)$$

$$\mathcal{G}^I(\theta) = \mathcal{D}_a E^{aI}(\theta) = \partial_a E^{aI}(\theta) + f^{IJK} A_{aJ}(\theta) E_K^a(\theta) \quad (4.54)$$

That is we have an infinite number of copies of the conventional complex $so(3)$ Ashtekar formulation, without any reality conditions. An infinite number of copies of complex gravity. The copies are labeled by the continuous parameter θ . Thus, from here it is clear that we really have an infinite number of degrees of freedom per point in spacetime; or we have four degrees of freedom per point in $\mathcal{M} \times S^1$. This means that if we choose to see the space S^1 as being a compactified fifth dimension, we really just have four degrees of freedom per point in the five-dimensional spacetime. On the other hand, if we insist on regarding the space S^1 as an abstract internal space, just introduced for notational convenience, we get an infinite number of degrees of freedom per point in spacetime.

If we want to select real general relativity, we know from the conventional Ashtekar formulation that we should have

$$\begin{aligned} Im(E^{aI}(\theta) E_I^b(\theta)) &= 0, & Im(E^{cI}(\theta) E^{(aJ}(\theta) \mathcal{D}_c E^{b)K}(\theta) f_{IJK}) &= 0 \\ Im(N(\theta)) &= 0, & Im(N^a(\theta)) &= 0 \end{aligned} \quad (4.55)$$

which may be translated back to the real $\widetilde{so(3)}$ formulation as

$$\begin{aligned} f^{nm}{}_k E_n^{aI} E_{Im}^b &= \delta_k^0 E_n^{aI} E_I^{bn}, & N_{(k)} &= N_{(-k)}, & N_{(k)}^a &= N_{(-k)}^a \\ f^{nm}{}_k f^{kl}{}_j E_m^{cI} E_l^{(aJ} \mathcal{D}_c E^{b)Kj} f_{IJK} &= \delta_0^n f^{kl}{}_j E_k^{cI} E_l^{(aJ} \mathcal{D}_c E^{b)Kj} f_{IJK} \end{aligned} \quad (4.56)$$

The field configuration (4.30)-(4.34) satisfies this.

If one is interested in knowing what the field configuration (4.30)-(4.34) corresponds to here in the S^1 -formulation, it is straightforward to Fourier expand it:

$$E^{aI}(\theta) = iE^{aI}f(\theta), \quad A_{aI}(\theta) = \Gamma_{aI}(E^{bJ}) - iK_{aI}f(\theta) \quad (4.57)$$

$$N(\theta) = N, \quad N^a(\theta) = N^a \quad (4.58)$$

where

$$f(\theta) = \begin{cases} 1, & 0 < \theta < \pi \\ -1, & \pi < \theta < 2\pi \end{cases} \quad (4.59)$$

Before leaving this theory, I want to comment on how to construct a related theory with a finite number of degrees of freedom per point. As we have seen above, the loop algebra $\widetilde{so(3)}$ is isomorphic to a direct sum of an infinite number of $so(3)$ algebras. What if we just consider a direct sum of N $so(3)$ algebras? Or put differently, the loop algebra can be seen as a map from S^1 to the base algebra. Alternatively we could consider a periodic one-dimensional lattice with N lattice points, and construct an "lattice-algebra" as a map from the lattice to the base-algebra. Explicitly one could start from N $so(3)$ valued fields $A_{aI}(n)$, $E^{aI}(n)$, $N(n)$, $N^a(n)$ and $A_{0I}(n)$ labeled by the integer n , and make the lattice periodic by requiring $E^{aI}(n+N) = E^{aI}(n)$ and similarly for the other fields. The next step to reach the "lattice-algebra" is to Fourier expand the fields:

$$E^{aI}(n) = \sum_{k=1}^N E_k^{aI} \exp(-ik\frac{2\pi n}{N}) \Rightarrow E_k^{aI} = \frac{1}{N} \sum_{n=1}^N E^{aI}(n) \exp(ik\frac{2\pi n}{N}) \quad (4.60)$$

and similarly for all the other fields. In the "lattice-algebra" formulation everything will look exactly as in the loop-algebra formulation in subsection 4.2 except that we now should identify two fields that differ by N steps in the level-index: $E_{k+N}^{aI} = E_k^{aI}$ etc. Then, we could again try to find the theory of Einstein gravity embedded in the *real* "lattice-algebra" theory, and it is easily shown that for N odd and > 1 , the following field configuration will work:

$$E^{aI}(n) = iE^{aI}f(n), \quad A_{aI}(n) = \Gamma_{aI}(E^{bJ}) - iK_{aI}f(n) \quad (4.61)$$

$$N(n) = N, \quad N^a(n) = N^a \quad (4.62)$$

where

$$f(n) = \begin{cases} 1, & 1 \leq n \leq \frac{N}{2} - 1 \\ -1, & \frac{N}{2} + 1 \leq n \leq N \end{cases} \quad (4.63)$$

and the Fourier transformed field E_k^{aI} etc. are given by (4.60).

5 Conclusions

The fact that the $so(1, 3; R)$ and the $so(\widetilde{3}; R)$ theory are shown to be equivalent to the $so(3; C)$ Ashtekar formulation, and infinitely many copies of it, should not come as a total surprise. That is just a reflection of the isomorphisms

$$so(1, 3; R) \cong so(3; C) \quad \text{and} \quad so(\widetilde{3}; R) \cong \sum_{n=-\infty}^{\infty} \oplus so_n(3; C). \quad (5.1)$$

The interesting thing about this formulations is that the reality conditions are transformed into normal constraint-like conditions on the fields. Thus, if it is possible to keep the reality conditions "outside" the theory, and only impose them in the quantum inner product³, then the same is possibly true for certain types of (second class) constraints. If that is the case, what is the criteria on the constraints, making this feasible? Guided by the real formulations in these paper, one may make the following conjecture: if the real theory has a complex structure, and the constraints are equivalent to requiring some objects to be "real" w.r.t this complex structure, then it ought to be possible to use these constraints to select an inner product.

More specifically: Take a real vector space $\mathcal{V}$ equipped with the complex structure J , such that $J^2 = -\mathbf{1}$, and a "complex conjugation" operator C , such that $C^2 = \mathbf{1}$ and $CJC = -J$. "Complex conjugation" is defined as $\bar{A} := CA$ for all vectors A in $\mathcal{V}$. An object – vector or function of vectors – is said to be "real" if it is invariant under C . An "imaginary" object changes sign under C . If we now, in our theory, have a set of constraints that is equivalent to requiring some objects to be "real", then we may put these constraints outside the theory, and later use them to select an inner product. The criteria we should have on the inner product is that the operators that correspond to the "real" objects, should be self adjoint.

In the real theories studied in this paper, the above objects may be specified as follows. For the $so(1, 3; R)$ theory: the vector space $\mathcal{V}$ is the space of $so(1, 3; R)$ valued functions on spacetime. The "real" subspace $\mathcal{V}_R$ is given by pure rotations, while $\mathcal{V}_I$ contains only pure boosts. "Complex conjugation" thus corresponds to changing sign of the boost-part of a general element in $\mathcal{V}$, and, using the basis (3.1), we may represent C as: $C = \text{diag}(-1, -1, -1, 1, 1, 1)$. The complex structure J is given by g_I^{*J} . It satisfies $g_I^{*J} g_J^{*K} = -\delta_I^K$ and it maps rotations into boosts and vice versa. A "real" object is invariant under "complex conjugation", and it is easy to check that *e.g.* $E^{aI} E_I^b$ is "real" while $E^{*aI} E_I^b$ is "imaginary".

In the $so(\widetilde{3}; R)$ theory, the vector space $\mathcal{V}$ is the space of $so(\widetilde{3}; R)$ valued functions on spacetime. The "real" subspace $\mathcal{V}_R$ consists of elements that is even under $n \rightarrow -n$, while the imaginary subspace $\mathcal{V}_I$ contains the odd elements. Here, n is the infinite dimensional level-index in $so(\widetilde{3}; R)$. The complex structure J is given by $J_n^m := f_n^{mk} g_k$, where g_k is the fourier coefficients of the imaginary step-function (4.30). This J satisfies $J_n^m J_m^k = -\delta_n^k$ and it maps even elements into odd ones, and vice versa. "Complex conjugation" is given by the simple operation $n \rightarrow -n$, and may be represented as: $C_n^m = \delta(n + m)$. Thus, we see that *e.g.* $E_n^{aI} E_I^{bn}$ is real, while $E_{(k)n}^{aI} E_I^{bn}$ is "complex".

³This is, so far, the standard way of postponing the problem of dealing with the reality conditions in the quantization program of the Ashtekar variables. One assumes that the reality conditions may be used to select the correct inner product.

Note that, for both these two examples, "complex conjugation" is really an automorphism of the real Lie-algebras $so(1, 3; R)$ and $so(3; R)$.

This shows that the constraints (3.56) and (4.56) can be interpreted as "reality conditions", and may thus possibly be used to select an inner product.

Note, though, that it is not yet clear that the reality conditions, in Ashtekar variables, can be successfully used to select an inner product. No one has yet shown how this should be done, in practice. I just say that *if* that is the case, then one should also be able to handle certain types of second class constraints the same way. (Actually, there is no reason to just restrict this to constraints of second class; even first class constraints might be handled this way.)

Finally, I want to comment on my starting motivation for studying these theories; I really looked for a generalization of Ashtekar's variables that could serve as a unified theory of gravity and Yang-Mills theory. The reason why the theories studied here only give complex gravity, may be understood from the fact that both $so(1, 3; R)$ as well as $so(3)$ are isomorphic to direct sums of $so(3; C)$ algebras. Therefore, to find something more non-trivial, one needs to find a Lie-algebra whose structure constants satisfy an identity of the requested type (2.5), but which is not isomorphic to sums of $so(3; C)$ algebras. I do not know of any such algebras. Also, although the theories described in this paper were shown to be equivalent to complex gravity, it is not clear what kind of linearized dynamics that would follow from *e.g.* a linearization of the real $so(1, 3)$ theory around de Sitter spacetime. It would indeed be interesting if for instance the $su(2)$ Yang-Mills theory falls out from the linearized theory. (Since the number of degrees of freedom in the $so(1, 3)$ theory is only four per spacetime point, the de Sitter solution has to be a very degenerate point in the space of solutions, for this to be possible.)

Acknowledgements

I thank Abhay Ashtekar, Fernando Barbero, Ingemar Bengtsson and Madhavan Varadarajan for interesting discussions. This work was supported by the NFR (Sweden) contract no. F-PD 10070-300.

References

- [1] A. Ashtekar, Phys. Rev. Lett. **57** (1986) 2244, Phys. Rev. **D36** (1987) 1587, A. Ashtekar, "Lectures on non-perturbative canonical gravity." (World Scientific, Singapore, 1991).
- [2] P. Peldán, Nucl. Phys. **B395** (1993) 239.
- [3] P. Peldán, Phys. Rev. D **46** (1992) R2279.
- [4] S. Chakraborty and P. Peldán, "Gravity and Yang-Mills theory: two faces of the same theory?" Penn State preprint CGPG-94/2-4, gr-qc/9403002, To appear in Int. J. Mod. Phys. D.
- [5] J. Samuel, Pramana-J Phys. **28** (1987) L429.

- [6] T. Jacobson and L. Smolin, *Class. Quant. Grav.* **5** (1988) 583.
- [7] R. Capovilla, J. Dell and T. Jacobson, *Class. Quant. Grav.* **8** (1991) 59.
R. Capovilla and T. Jacobson, *Class. Quant. Grav.* **9** (1992) 1839.
R. Capovilla, J. Dell and T. Jacobson, *Phys. Rev. Lett.* **63** (1989) 2325
R. Capovilla, J. Dell, T. Jacobson, and L. Mason, *Class. Quant. Grav.* **8** (1991) 41.
- [8] P. Peldán, *Class. Quant. Grav.* **8** (1991) 1765.
- [9] P. Peldán, "Actions for gravity, with generalizations: A review", gr-qc/9305011, To appear in *Class. Quant. Grav.*
- [10] A. Pressley and G. Segal, "Loop Groups", (Oxford University Press, New York, 1986)
P. Goddard and D. Olive, *Int. J. Mod. Phys.* **A1** (1986) 303. (1976) 88.